



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

No. IV.

A short and easy rule for finding the equation for the change of the sun's declination when equal altitudes are used to regulate a clock or other time keeper. Communicated by Andrew Ellicott Esq.

Read January 16th, 1801.

FOR THE FIRST PART.

FIND the Sun's longitude, declination, and the change of declination for 24^h at the time of the observation, likewise find the proportional part of the change of declination for the half interval between the forenoon and afternoon observations, then take the proportional logarithm answering to the change of declination for the half interval, (increasing the index by 10,) from which take the log. cosecant of the horary angle; to the remainder add the log. cotangent of the latitude of the place of observation, and take out the minute and second from the P. Ls. answering to the sum (10 being deducted from the index) which converted into time will give the first part of the correction and will be deductive in North latitudes, when the sun's longitude is 0, 1, 2, 9, 10, or 11, signs, and additive in the others; but the contrary in South latitudes.

FOR THE SECOND PART.

TO the P. L. of the change of the sun's declination during the half interval, add the log. cotangent of the sun's declination, from that sum deduct the log. cotangent of the horary angle.—Take out the minute and second from P. Ls. answering to the remainder, which turned into time will give the second part of the correction; this is common to all latitudes, and will be additive when the sun's longitude is 0, 1, 2, 6, 7, or 8, signs, and deductive in the others.

EXAMPLE.

*Suppose the following equal altitudes were taken in latitude $39^{\circ}.56'$.
N. when the sun's longitude was $4^s. 15^{\circ}$.*

A. M. $8^h. 32' 20''$ —P. M.	$3^h 32' 24''$
Add.....	<u>12 0 0</u>
	15 32 24
Deduct forenoon's observation.....	<u>8 32 20</u>
	2) 7 0 4
half interval.....	<u>3 30 2</u>
Add forenoon's observation.....	<u>8 32 20</u>
Sun's center on the meridian nearly	<u><u>12 2 22</u></u>

FOR THE CORRECTION.

The sun's declination answering to $4^s 15^{\circ}$ of his longitude is nearly $16^{\circ} 21'$, and the change of declination at the same time about $16' 55''$ in 24 hours, or $2' 28''$ during the half interval.

THEN BY THE RULE.

Change of declination during

half interval $2' 28''$ P. L. 11. 8631.

Horary angle $52^{\circ} 30'$ log. cosec.—10. 1005

1. 7626

Latitude $39^{\circ} 56'$ log. cotan. +10. 0772

P. L. 1. 8398 = $2' 36'' = 10'' 24'''$ in

time, being the first part of the equation, and additive by the rule.

FOR THE SECOND PART.

Change of declination during the	
half interval 2' 28" P. L.	1. 8631
Sun's declination 16° 21' log cotan. +	10. 5326
	<u>12. 3957</u>
Horary angle 52' 30' log. cotan. —	9. 8850
P. L. —	<u>2. 5107 = 0' 33" = 2" 12" in</u>

time, being the second part of the equation, and deductive by the rule,

APPLICATION.

Apparent time of the sun's center on } the meridian by equal altitudes nearly }	12 ^h 2' 22" 0 ^m
First part of the equation	+ 10 ^m 24 ^m
Second do.....—	2. 12 + 8. 12
Sun's centre on the meridian	<u>12. 2. 30. 12.</u>

No. V.

Account of an extraordinary flight of meteors (commonly called shooting of stars) communicated by Andrew Ellicot, Esq. as extracted from his Journal in a voyage from New-Orleans to Philadelphia.

Read 16th January, 1801

“ NOVEMBER 12th 1799, about three o'clock, A. M. I was called up to see the shooting of the stars (as it is commonly called.) The phenomenon was grand and awful, the whole heavens appeared as if illuminated with sky-rockets, which disappeared only by the light of the sun after day break. The meteors, which at any one instant of time appeared as nume-